


Recurrent Urinary Tract Infections in Patients with Type 2 Diabetes Mellitus: A Narrative Review

Maher Kadhum Abed^{1*}, Manal Khalid Abdulridha², Hussein Adnan³

¹ Department of Clinical Pharmacy, College of Pharmacy, Al-Mustansiriyah University, Baghdad, Iraq

² Department of Clinical Pharmacy, College of Pharmacy, Al-Mustansiriyah University, Baghdad, Iraq

³ Department of Internal Medicine, College of Medicine, Wasit University, Wasit, Iraq

 <https://doi.org/10.60988/p.v38i2.305>

KEYWORDS: recurrent urinary tract infection; type 2 diabetes mellitus; antimicrobial resistance; probiotics; SGLT2 inhibitors

ARTICLE INFO:

Received: January 13, 2026

Revised: January 28, 2026

Accepted: February 23, 2026

Available on line: May 17, 2026

***CORRESPONDING AUTHOR:**

Maher Kadhum Abed E-mail:

maherkadhum@uomustansiriyah.edu.iq

mahir.kadhim@gmail.com

ABSTRACT

Recurrent urinary tract infections (rUTIs) represent a frequent and clinically challenging complication among patients with type 2 diabetes mellitus (T2DM). Impaired glycemic control alters host defense mechanisms, promotes glycosuria, and facilitates bacterial adhesion and colonization within the urinary tract, thereby increasing susceptibility to recurrent infections. This narrative review summarizes current evidence regarding the epidemiology, pathophysiology, microbial characteristics, diagnostic considerations, and management strategies of rUTIs in individuals with T2DM. *Escherichia coli* remains the predominant uropathogen; however, diabetic patients are disproportionately affected by multidrug-resistant organisms, complicating therapeutic decision-making. Autonomic neuropathy and bladder dysfunction further contribute to urinary stasis and infection recurrence. Diagnosis relies on careful clinical assessment supported by urine microscopy and culture, which is particularly important in diabetic patients to guide appropriate antimicrobial therapy. Management principles are generally similar to those applied in non-diabetic individuals but require classification of most infections as complicated, emphasizing culture-guided antibiotic selection, appropriate treatment duration, and vigilant monitoring. Non-pharmacological measures, strict glycemic control, and evaluation for structural or functional urinary tract abnormalities are essential components of care. Emerging therapies, including sodium-glucose cotransporter-2 inhibitors and non-antibiotic preventive approaches such as probiotics, warrant further investigation. An integrated, multidisciplinary approach combining optimized diabetes management with judicious antimicrobial use is critical to reducing recurrence, preventing complications, and limiting antimicrobial resistance in this vulnerable population.

1. Introduction

Urinary tract infections (UTIs) are among the most common bacterial infections encountered in clinical practice. Patients with T2DM exhibit a higher incidence of both uncomplicated and complicated UTIs compared with non-diabetic individuals¹. Chronic hyperglycemia contributes to immune dysfunction, microvascular damage, and autonomic neuropathy, all of which predispose diabetic patients to recurrent urinary tract infections (rUTIs). The pathogenesis of rUTI in diabetic individuals may be caused by a number of immune system abnormalities, poor metabolic control of diabetes, and incomplete bladder evacuation brought on by autonomic neuropathy². Diabetes alters the natural host system, which may contribute to the development of UTIs³. Limited research indicates that defective bladder contractions may result in stagnant urine pools, which provide a conducive environment for bacterial proliferation. Additional research suggests that urine with elevated glucose levels fosters a rise in bacterial clusters and their colonization within the urinary system⁴. Concerning the findings of therapeutic approaches for asymptomatic bacteriuria in diabetic individuals, limited clinical trials have been performed. The findings of these clinical trials demonstrate that treatment durations of 2 weeks and 6 weeks exhibit equivalent efficacy, the incidence of rUTIs remains elevated, despite prolonged antibiotic therapy, the majority of re-infections occurred recurrently (4/8 weeks post-therapy) and these recurrences were not caused by the same microorganism. Physicians should be cognizant of the elevated incidence of underlying structural genitourinary abnormalities in bacteriuric women with diabetes mellitus⁵. In addition to increasing infection risk, diabetes is associated with more severe clinical presentations and higher rates of complications, including emphysematous cystitis and pyelonephritis⁶. This review provides an updated narrative overview of the mechanisms, diagnosis, and management of rUTIs in patients with T2DM.

2. Pathophysiology of rUTIs in T2DM

2.1 Impaired host defense mechanisms

In diabetic patients, the elevated risk of rUTI may be attributed to a variety of potential mechanisms that are distinctive to diabetes⁵. The proliferation of pathogenic bacteria may be facilitated by elevated glucose concentrations in urine⁷. Elevated renal parenchymal glucose concentrations foster an environment conducive to the proliferation of bacteria, potentially serving as a precipitating factor for pyelonephritis and renal consequences, including emphysematous pyelonephritis⁸. Various abnormalities in the immune system, including humoral, cellular, and innate immunity, may contribute to the pathogenesis of infections of the urinary system in patients with diabetes⁹. Patients with diabetes and asymptomatic bacteriuria exhibited reduced levels of interleukin-6 and interleukin -8 compared to their non-diabetic counterparts with asymptomatic bacteriuria¹⁰. Autonomic neuropathy affecting the genitourinary system leads to impaired voiding and urine retention, which diminishes physical bacterial elimination by urination, consequently promoting bacterial proliferation¹¹. Bladder dysfunction impacts 26%–85% of diabetic women, depending on age, severity of neuropathy, and duration of diabetes¹². The emergence of new anti-diabetic sodium glucose cotransporter-2 inhibitors SGLT2 inhibitors, which promote glycosuria, has elicited worries about a possible increase in urinary tract infections¹³. Thus, the most frequent side effect of SGLT2 inhibitors therapy seems to be genitourinary tract infections. Increased urinary glucose can lead to a faster rate of microbial development, which raises the risk of ASB and UTI. Urine that is subsequently deposited on the vulvovaginal tissues after voiding can further raise the risk of urogenital infections¹⁴. Non-DM individuals with difficult UTIs had similar microorganisms to those recovered from DM patients with UTIs. Enterobacter spp., Proteus spp., E. coli, Klebsiella spp., Group B Streptococci, and Enterococcus faecalis are the most commonly isolated pathogens¹⁵. Figure 1 summarizes the steps




Figure 1. The pathophysiology of urinary tract infections. ,

that contribute to the pathophysiology of UTIs.

2.2 Glycosuria and bacterial proliferation

Persistent hyperglycemia results in glycosuria, creating a favorable environment for bacterial growth within the urinary tract. Elevated glucose concentrations in renal parenchyma and urine enhance bacterial adhesion and colonization, increasing the risk of ascending infections and severe renal complications¹⁶.

2.3 Bladder dysfunction and urinary stasis

[Diabetic autonomic neuropathy affects bladder sensation and contractility, leading to incomplete bladder emptying and increased post-void residual volume. Bladder dysfunction has been reported in up to 87% of diabetic women and represents a key contributor to urinary stasis and recurrent infection¹⁷.

3. Microbiological Patterns

The spectrum of uropathogens isolated from diabetic patients with rUTIs is broadly similar to that observed in non-diabetic populations. *Escherichia coli* remains the most frequently identified organism, followed by *Klebsiella* species, *Proteus* species, *Enterococcus faecalis*, and Group B *Streptococci*¹⁸.

4. Diagnosis of Recurrent UTIs in T2DM

4.1 Clinical features

The clinical manifestation of rUTI is identical to that of sporadic acute cystitis. Local genitourinary symptoms of dysuria, frequency, urgency, and hesitation manifest abruptly¹⁹). In addition, simple cystitis may show with suprapubic discomfort and gross hematuria. The symptoms are frequently the same as in earlier episodes, and ambulatory women with recurrent, simple UTIs are quite accurate in diagnosing themselves¹⁹

A UTI diagnosis should be taken into consideration

for any diabetic patient displaying symptoms suggestive of a UTI. According to a recent multi-center study on women with community-acquired acute pyelonephritis in South Korea, diabetic patients experienced significantly fewer episodes of lower urinary tract infection symptoms, costovertebral angle soreness, and flank pain than their non-diabetic counterparts²⁰. Patients with T2DM and UTIs may exhibit hypo- or hyperglycemia, non-ketotic hyperosmolar condition, or ketoacidosis, necessitating the immediate elimination of infectious triggering factors, including UTIs²¹.

4.2 Laboratory investigations

Urinalysis typically demonstrates pyuria, which can be detected by microscopy or leukocyte esterase testing. Urine culture is strongly recommended prior to initiating antimicrobial therapy in diabetic patients to distinguish infection from colonization and to guide antibiotic selection. In recurrent cases, imaging studies may be required to evaluate for underlying structural abnormalities²².

5. Management Strategies

5.1 Non-pharmacological management

General preventive measures include adequate hydration, regular and complete bladder emptying, avoidance of unnecessary catheterization, and modification of behavioral risk factors such as spermicide use. Cranberry products and probiotics have demonstrated modest preventive benefits in selected populations, although evidence specific to diabetic patients remains limited²³.

5.2 Pharmacological management

Antimicrobial medications are frequently used to treat symptomatic UTIs. Individualized treatment should be provided based on the antibiotic sensitivity of the urine culture, or a broad-spectrum antimicrobial medication may be used as empirical or preventive therapy²⁴. Commonly utilized antibiotics

comprise trimethoprim/sulfamethoxazole, cotrimoxazole, amoxicillin, quinolones, and nitrofurantoin²⁵. For people with acute uncomplicated cystitis, the first-line treatment is oral nitrofurantoin for five days; fosfomycin is an alternate. These two medications, however, do not work well on renal parenchyma and are therefore not advised for pyelonephritis; instead, ciprofloxacin and trimethoprim/sulfamethoxazole are utilized²⁶.

people with diabetes who have acute pyelonephritis are more likely to experience more serious consequences, such as bacteremia, extended hospitalization, or death, than people without diabetes.²⁷ Bilateral acute pyelonephritis occurs more frequently in individuals with T2DM²⁸. Management of pyelonephritis in diabetic patients presenting with pronounced symptoms—such as nausea, vomiting, or metabolic disturbances—requires initial administration of intravenous antibiotics during hospitalization, while those with milder symptoms can be treated with oral antibiotics²¹. Effective glycemic control, whether through insulin or other hypogly-

cemic agents, is a vital component of therapy²⁹. Furthermore, surgical interventions including drainage, debridement, or computed tomography (CT)-guided procedures may be necessary, depending on clinical severity³⁰. In critical presentations such as sepsis, broad-spectrum antibiotics must be administered urgently—ideally within the first hour—alongside fluid resuscitation and prompt diagnostic evaluation via blood and urine cultures.³¹.

6. Conclusions

Recurrent UTIs in patients with T2DM represent a multifactorial clinical challenge influenced by metabolic, immunological, and functional abnormalities. Effective management requires an integrated approach combining strict glycemic control, appropriate diagnostic evaluation, and judicious antimicrobial use. Future research should focus on individualized prevention strategies, optimization of non-antibiotic therapies, and the long-term impact of novel antidiabetic agents on urinary tract infection risk.

References

- Confederat L.G., Condurache M.I., Alexa R.E., Dragostin O.M. Particularities of urinary tract infections in diabetic patients: a concise review. *Medicina (B Aires)*. 59(10),1747,2023.
- Truzzi J. C. I., Almeida F. M. R., Nunes E. C., Sadi M. V. Residual Urinary Volume and Urinary Tract Infection-When are They Linked? *J Urol*. 180(1), 182–185,2008.
- Kumar Jha P, Baral R., Khanal B. Prevalence of uropathogens in diabetic patients and their susceptibility pattern at a tertiary care center in Nepal-a retrospective study. *Int J Bio Lab Sci*. 3, 29–34,2014.
- Chen S. L., Jackson S. L., Boyko E. J. Diabetes mellitus and urinary tract infection: epidemiology, pathogenesis and proposed studies in animal models. *J. Urol*. 182, S51–S56,2009.
- Akash M. S. H., Rehman K., Fiayyaz F., Sabir S., Khurshid M. Diabetes-associated infections: development of antimicrobial resistance and possible treatment strategies. *Arch. Microbiol*. 202, 953–965,2020.
- Casqueiro J., Casqueiro J., Alves C. Infections in patients with diabetes mellitus: A review of pathogenesis. *Indian J. Endocrinol. Metab*. 16, S27–S36,2012.
- Wang M.-C. *et al*. Bacterial characteristics and glycemic control in diabetic patients with Escherichia coli urinary tract infection. *JMII* 46, 24–29,2013.
- Schneeberger C. *et al*. Differences in the pattern of antibiotic prescription profile and recurrence rate for possible urinary tract infections in women with and without diabetes. *Diabetes Care*. 31, 1380–1385,2008.
- Geerlings S. E. *et al*. Cytokine secretion is impaired in women with diabetes mellitus. *Eur J Clin Invest*. 30, 995–1001,2000.
- Soo Park B., *et al*. Outcome of nephrectomy and kidney-preserving procedures for the treatment of emphysematous pyelonephritis. *Scand J Urol*.

- Nephrol.* 40, 332–338, 2006.
11. Truzzi J. C. I., Almeida F. M. R., Nunes E. C. , Sadi M. V. Residual urinary volume and urinary tract infection—when are they linked? *J Urol.* 180, 182–185, 2008.
 12. Nitzan O., Elias M., Chazan B. & Saliba W. Urinary tract infections in patients with type 2 diabetes mellitus: review of prevalence, diagnosis, and management. *Diabetes Metab Syndr Obes.* 129–136, 2015.
 13. Ptaszynska A., et al. Safety Profile of Dapagliflozin for Type 2 Diabetes: Pooled Analysis of Clinical Studies for Overall Safety and Rare Events. *Drug Saf.* 37, 815–829, 2014.
 14. Soh P. N., et al. Urinary and genital infections in patients with diabetes: how to diagnose and how to treat. *Diabetes Metab.* 42, 16–24, 2016.
 15. Hamdan H. Z., et al. Urinary tract infections and antimicrobial sensitivity among diabetic patients at Khartoum, Sudan. *Ann. Clin. Microbiol. Antimicrob.* 14, 26, 2015.
 16. John P. P., et al. Exposure to moderate glycosuria induces virulence of group B streptococcus. *J Infect. Dis.* 223, 843–847, 2021.
 17. Golbidi S. , Laher I. Bladder dysfunction in diabetes mellitus. *Front. Pharmacol.* 1, 136, 2010.
 18. Clare E. C., Juliet A. O., Maureen O. C., Romeo A. J., Ada K.O. E. Urinary Tract infections and antimicrobial sensitivity patterns of uropathogens isolated from diabetic and non-diabetic patients attending some hospitals in Awka. *Am. J. Microbiol. Res* 9, 83–91, 2021.
 19. Gupta K. , Trautner B. W. Diagnosis and management of recurrent urinary tract infections in non-pregnant women. *Bmj* 346, 2013.
 20. Kim Y. ,et al. Comparison of the clinical characteristics of diabetic and non-diabetic women with community-acquired acute pyelonephritis: a multicenter study. *J. Infect.* 69, 244–251, 2014.
 21. Nitzan O., Elias M., Chazan B. , Saliba W. Urinary tract infections in patients with type 2 diabetes mellitus: review of prevalence, diagnosis, and management. *Diabetes Metab. Syndr. Obes.* 129–136, 2015.
 22. Nicolle L. E. Urinary tract infection. *Crit. Care Clin.* 29, 699–715, 2013.
 23. Barea B. M., Veeratterapillay R. , Harding C. Non-antibiotic treatments for urinary cystitis: an update. *Curr. Opin. Urol.* 30, 845–852, 2020.
 24. George C. E., Norman G., Ramana G. V., Mukherjee D. , Rao T. Treatment of uncomplicated symptomatic urinary tract infections: Resistance patterns and misuse of antibiotics. *J. Family Med. Prim. Care* 4, 416–421, 2015.
 25. Bryce A., et al. Global prevalence of antibiotic resistance in paediatric urinary tract infections caused by Escherichia coli and association with routine use of antibiotics in primary care: systematic review and meta-analysis. *bmj* 352, 2016.
 26. Al Lawati H., Blair B. M. , Larnard J. Urinary tract infections: core curriculum 2024 *AJKD.* 83, 90–100, 2024.
 27. Kofteridis D. P., et al. Effect of diabetes mellitus on the clinical and microbiological features of hospitalized elderly patients with acute pyelonephritis. *J. Am. Geriatr. Soc.* 57, 2125–2128, 2009.
 28. Mogensen C. E. *The Kidney and Hypertension in Diabetes Mellitus.* Springer, 2013.
 29. Ronald A. , Ludwig E. Urinary tract infections in adults with diabetes. *Int. J. Antimicrob. Agents* 17, 287–292, 2001.
 30. Kamei J. , Yamamoto S. Complicated urinary tract infections with diabetes mellitus. *Journal of Infection and Chemotherapy* 27, 1131–1136, 2021.
 31. Levy M. M., Evans L. E. , Rhodes, A. The surviving sepsis campaign bundle: 2018 update. *Intensive Care Med.* 44, 925–928, 2018.